

Depression

References

- Agency for Healthcare Research and Quality.(2002). *Management of cancer symptoms: Pain, depression, and fatigue*. Evidenced Report/Technology Assessment: Number 61 (AHRQ Publication No. 02-E031). Rockville, MD.
- American Psychiatric Association. (2000, April). *Practice guideline for the treatment of patients with major depressive disorder* (2nd ed.). Retrieved November 10, 2006, from http://www.psych.org/psych_pract/treatg/pg/Practice%20Guidelines8904/MajorDepressiveDisorder_2e.pdf
- Antoni, M.H., Lehman, J.M., Kilbourn, K.M. Boyers, A.E. Culver, J.L., Alferi, S.M. et al. (2001). Cognitive-behavioral stress management intervention decreases the prevalence of depression and enhances benefit finding among women under treatment for early-stage breast cancer. *Health Psychology*, 20 (1), 20-32.
- Barsevick, A.M., Sweeney, C., Haney, E., & Chung, E. (2002). A systematic qualitative analysis of psychoeducational interventions for depression in patients with cancer. *Oncology Nursing Forum*, 29, 73–84.
- Bennett, G., & Badger, T.A. (2005). Depression in men with prostate cancer. *Oncology Nursing Forum*, 32, 545–556.
- Berglund, G., Petersson, L-M, Eriksson, K.C., Wallenius, I., Roshanai, A. Nordin, K.M., et al. (2007). “Between Men”: A psychosocial rehabilitation programme for men with prostate cancer. *Acta Oncological*, 46, 83-89.
- Cassileth, B.R., & Vickers, A.J. (2004). Massage therapy for symptom control: Outcome study at a major cancer center. *Journal of Pain and Symptom Management*, 28, 244–249.
- Davis, M., Eshelman, E.R., & McKay, M. (2000). *The relaxation and stress reduction workbook* (5th ed.). Oakland, CA: New Harbinger Publications, Inc.
- Fellowes, D., Barnes, K., & Wilkinson, S. (2004). Aromatherapy and massage for symptom relief in patients with cancer. *Cochrane Database of Systematic Reviews* 2004, Issue 3. Art. No.: CD002287.
- Given, C., Given, B., Rahbar, M., Jean, S., McCorkle, R., & Cimprich, B. (2004). Does a symptom management intervention affect depression among cancer patients: Results from a clinical trial. *Psycho-Oncology*, 13, 818–830.
- Goodnick, P.J., & Hernandez, M. (2000). Treatment of depression in comorbid medical illness. *Expert Opinion on Pharmacotherapy*, 1, 1367–1384.
- Gotay, C.C., Moinpour, C.M., Unger, J.M., Jiang, C.S., Coleman, D., Martino, S., et al. (2007). Impact of a peer-delivered telephone intervention for women experiencing a breast cancer recurrence. *Journal of Clinical Oncology*, 25 (15), 2093-2098.
- Hanna, L.R., Avila, P.F., Meter, J.D., Nicholas, D.R. & Kaminsky, L.A. (2008). The effects of a comprehensive exercise program on physical function, fatigue, and mood in patients with various types of cancer. *Oncology Nursing Forum*, 35 (3), 461-469.
- Helgeson, V.S., Lepore, S.J., & Eton, D.T. (2006). Moderators of the benefits of psychoeducational interventions for men with prostate cancer. *Health Psychology*, 25 (3), 348–354.

Depression

References

- Hernandez-Reif, M., Ironson, G., Field, T., Hurley, J., Katz, G., Diego, M., et al. (2004). Breast cancer patients have improved immune and neuroendocrine functions following massage therapy. *Journal of Psychosomatic Research*, *57*, 45–52.
- Homsy, J., Nelson, K.A., Sarhill, N., Rybicki, L., LeGrand, S.B., Davis, M.P., et al. (2001). A phase II study of methylphenidate for depression in advanced cancer. *American Journal of Hospice and Palliative Care*, *18*, 403–407.
- Lan Ly, K.L., Chidgey, J., Addington-Hall, J., & Hotopf, M. (2002). Depression in palliative care: A systematic review. Part 2: Treatment. *Palliative Medicine*, *16*, 279–284.
- Lewis, F.M., Casey, S.M., Brandt, P.A., Shands, M.E., & Zahlis, E.H. (2006). The enhancing connections program: Pilot study of a cognitive-behavioral intervention for mothers and children affected by breast cancer. *Psycho-Oncology*, *15*(6), 486–497.
- Luebbert, K., Dahme, B., & Hasenbring, M. (2001). The effectiveness of relaxation training in reducing treatment related symptoms and improving emotional adjustment in acute and non-surgical cancer treatment: A meta-analytical review. *Psycho-Oncology*, *10*, 490–502.
- Manne, S.L., Rubin, S., Edelson, M., Rosenblum, N., Bergman, C., Hernandez, E., et al. (2007). Coping and communication-enhancing intervention versus supportive counseling for women diagnosed with gynecological cancers. *Journal of Consulting and Clinical Psychology*, *75* (4), 615–628.
- Midtgaard, J., Rorth, M., Stelter, R., Tveteras, A., Andersen, C., Quist, M., et al. (2005). The impact of a multidimensional exercise program on self-reported anxiety and depression in cancer patients undergoing chemotherapy: A phase II study. *Palliative and Supportive Care*, *3*(3), 197–208.
- Ministry of Health, Singapore. (2004). *Clinical practice guidelines: Depression*. Retrieved November 10, 2006, from http://www.moh.gov.sg/corp/publications/list.do?id=pub_guide_clinics
- Musselman, D.L., Lawson, D.H., Gumnick, J.F., Manatunga, A.K., Penna, S., Goodkin, R.S., et al. (2001). Paroxetine for the prevention of depression induced by high-dose interferon alfa. *New England Journal of Medicine*, *344*, 961–966.
- National Cancer Institute. (2006). *Depression PDQ®* (Health Professional Version). Retrieved November 10, 2006, from http://cancer.gov/cancer_information/coping.
- National Comprehensive Cancer Network. (2006). *Clinical practice guidelines in oncology: Distress management, version 1.2007*. Retrieved November 10, 2006, from http://www.nccn.org/professionals/physician_gls/PDF/distress.pdf
- National Health and Medical Research Council (Australia). (2003). *Clinical practice guidelines for the psychosocial care of adults with cancer*. Retrieved November 10, 2006, from <http://www.nhmrc.gov.au/publications/synopses/cp90syn.htm>
- Newell, S.A., Sanson-Fisher, R.W., & Savolainen, N.J. (2002). Systematic review of psychological therapies for cancer patients: Overview and recommendations for future research. *Journal of the National Cancer Institute*, *94*, 558–584.

Depression

References

- Osborn, R.L., Demoncada, A.C., & Feuerstein, M. (2006). Psychosocial interventions for depression, anxiety, and quality of life in cancer survivors: Meta-analyses. *International Journal of Psychiatry in Medicine, 36*, 13–34.
- Pirl, W.F. (2004). Evidence report on the occurrence, assessment, and treatment of depression in cancer patients. *Journal of the National Cancer Institute Monographs, 32*, 32–39.
- Pugliese, P., Perrone, M., Nisi, E., Garufi, C., Giannarelli, D., Bottomley, A., et al. (2006). An integrated psychological strategy for advanced colorectal cancer patients. *Health and Quality of Life Outcomes, 4*, 1–9.
- Qaseem, A., Snow, V., Shekelle, P., Casey, D.E., Cross, J.R., & Owens, D.K. (2008). Evidence-Based interventions to improve the palliative care of pain, dyspnea, and depression at the end of life: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine, 148*, 141-146.
- Rajasekaran, M., Edmonds, P.M., & Higginson, I.L. (2005). Systematic review of hypnotherapy for treating symptoms in terminally ill adult cancer patients. *Palliative Medicine, 19*, 418–426.
- Rodin, G., Lloyd, N., Katz, M., Green, E., Mackay, J.A., & Wong, R.K.S. (2007). The treatment of depression in cancer patients: A systematic review. *Supportive Care in Cancer, 15*, 123-136.
- Rozans, M., Dreisbach, A., Lertora, J.J., & Kahn, M.J. (2002). Palliative uses of methylphenidate in patients with cancer: A review. *Journal of Clinical Oncology, 20*, 335–339.
- Sadock, B.J., & Sadock, V.A. (2003). *Kaplan & Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry* (9th ed.). Philadelphia: Lippincott Williams and Wilkins.
- Seaward, B.L. (2004). *Managing stress: Principles and strategies for health and wellbeing* (4th ed.). Sudbury, MA: Jones and Bartlett Publishers.
- Scheier, M.F., Helgeson, V.S., Schulz, R., Colvin, S., Berga, S., Bridges, M.W., et al. (2005). Interventions to enhance physical and psychological functioning among younger women who are ending nonhormonal adjuvant treatment for early-stage breast cancer. *Journal of Clinical Oncology, 23* (19), 4298-4311.
- Schwartz, L., Lander, M., & Chochinov, H.M. (2002). Current management of depression in cancer patients. *Oncology, 16*, 1102–1115.
- Sharpe, M., Strong, V., Allen, K., Rush, R., Maguire, P., House, A., et al. (2004). Management of major depression in outpatients attending a centre: A preliminary evaluation of a multicomponent cancer nurse-delivered intervention. *British Journal of Cancer, 90*, 310–313.
- Sloman, R. (2002). Relaxation and imagery for anxiety and depression control in community patients with advanced cancer. *Cancer Nursing, 25*, 432–435.
- Soden, K., Vincent, K., Craske, S., Lucas, C. & Ashley, S. (2004). A randomized controlled trial of aromatherapy massage in a hospice setting. *Palliative Medicine, 18*, 87-92.



Depression

References

- Thastum, M., Munch-Hansen, A., Wiell, A., & Romer, G. (2006). Evaluation of a focused short-term preventive counseling project for families with a parent with cancer. *Clinical Child Psychology and Psychiatry, 11*(4), 529–542.
- Uitterhoeve, R.J., Vernooy, M., Litjens, M., Potting, K., Bensing, J., De Mulderf, P., et al. (2004). Psychosocial interventions for patients with advanced cancer—A systematic review of the literature. *British Journal of Cancer, 91*, 1050–1062.
- Vilela, L.D., Nicolau, B., Mahmud, S., Edgar, L., Hier, M., Black, M., et al. (2006). Comparison of psychosocial outcomes in head and neck cancer patients receiving a coping strategies intervention and control subjects receiving no intervention. *Journal of Otolaryngology, 35* (2), 88-96.
- Williams, S., & Dale, J. (2006). The effectiveness of treatment for depression/depressive symptoms in adults with cancer: A systematic review. *British Journal of Cancer, 94*, 372–390.