

Lymphedema References

- Ahmed, R.L., Thomas, W., Yee, D., & Schmitz, K.H. (2006). Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *Journal of Clinical Oncology*, 24(18), 2765-2772.
- Badger, C., Preston, N., Seers, K. & Mortimer, P. (2004). Physical therapies for reducing and controlling lymphoedema of the limbs. *Cochran Database of Systematic Reviews*, 4, CD003141.
- Bani, H.A., Fasching, P.A., Lux, M.M., Rauh, C., Willner, M., Eder, I., et al. (2007). Lymphedema in breast cancer survivors: Assessment and information provision in a specialized breast unit. *Patient Education and Counseling*, 66(3), 311-318.
- Barclay, J., Vestey, J., Lambert, A., & Balmer, C. (2006). Reducing the symptoms of lymphoedema: Is there a role for aromatherapy? *European Journal of Oncology Nursing*, 10(2), 140-149.
- Bernard, P. (2008). Management of common bacterial infections of the skin. *Current Opinion in Infectious Diseases*, 21(2), 122-128.
- Bicego, D., Brown, K., Ruddick, M., Storey, D., Wong, C., & Harris, S.R. (2006). Exercise for women with or at risk for breast cancer-related lymphedema. *Physical Therapy*, 86(10), 1398-1405.
- Box, R.C., Reul-Hirsch, H.M., Bullock-Saxton, J.E., & Furnival, C.M. (2002). Physiotherapy after breast cancer surgery: Results of a randomised controlled study to minimise lymphoedema. *Breast Cancer Research and Treatment*, 75(1), 51-64.
- British Lymphology Society, (1999) *Guidelines for the use of MLD and SLD*. Retrieved November 21, 2007 from <http://www.lymphoedema.org/bls/membership/guidelines.htm>.
- Carati, C.J., Anderson, S.N., Gannon, B.J., & Piller, N.B. (2003). Treatment of post-mastectomy lymphedema with low-level laser therapy. *Cancer*, 98(6), 1114-1122.
- Casley-Smith, J.R. & Casley-Smith, J.R. (1996). Lymphedema initiated by aircraft flights. *Aviation, Space and Environmental Medicine*, 67 (1), 52-56.
- Centers for Disease Control and Prevention. (2008). *Body mass index*. Retrieved July 31, 2008 from <http://www.cdc.gov/nccdphp/dnpa/healthyweight/assessing/bmi/index.htm>.
- Chapman, D.D. (2007). Local and regional contraol. In S.M. Mahon (Ed.) *Site-specific cancer series:Breast cancer* (pp. 63-80). Pittsburgh, PA., Oncology Nursing Society.
- Chapman, D.D. & Moore, S. (2005) Breast cancer in C.H. Yarbrow, M.H. Frogge & M. Goodman (Eds.), *Cancer nursing: Principles and practice* (6th ed., pp.1022-1088). Sudbury, MA: Jones and Bartlett
- Cheema, B., Gaul, C.A., Lane, K., & Fiatarone Singh, M.A. (2008). Progressive resistance training in breast cancer: A systematic review of clinical trials. *Breast Cancer Research and Treatment*, 109(1), 9-26.
- Cinar, N., Seckin, U., Kreskin, D., Bodur, H., Bozkurt, B., & Cengiz, O. (2008). The effectiveness of early rehabilitation in patients with modified radical mastectomy. *Cancer Nursing*, 31(2), 160-165.
- de Rezende, L.F., Franco, R.L., de Rezende, M.F., Beletti, P.O., Morais, S.S., & Gurgel, M.S. (2006). Two exercise schemes in postoperative breast cancer: Comparison of effects on shoulder movement and lymphatic disturbance. *Tumori*, 92(1), 55-61.
- Didem, K., Ufuk, Y.S., Serdar, S., & Zumre, A. (2005). The comparison of two different

- physiotherapy methods in treatment of lymphedema after breast surgery. *Breast Cancer Research and Treatment*, 93(1), 49-54.
- Forner-Cordero, I., Navarro-Monsolieu, R., Munoz-Langa, J., Alcober-Fuster, P., & Rel-Monzo, P. (2007). Use of a nanocrystalline silver dressing on lymphatic ulcers in patients with chronic lymphoedema. *Journal of Wound Care*, 16(5), 235-239.
- Gothard, L., Stanton, A., MacLaren, J., Lawrence, D., Hall, E., Mortimer, P., et al. (2004). Non-randomized phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphedema and tissue fibrosis after radiotherapy for early breast cancer. *Radiotherapy and Oncology*, 70(3), 217-224.
- Graham, P.H. (2002). Compression prophylaxis may increase the potential for flight-associated lymphoedema after breast cancer treatment. *Breast*, 11 (1), 66-71.
- Hamner, J.B., & Fleming, M.D. (2007). Lymphedema therapy reduces the volume of edema and pain in patients with breast cancer. *Annals of Surgical Oncology*, 14(6), 1904-1908.
- Indelicato, D.J., Grobmyer, S.R., Newlin, H., Morris, C.G., Haigh, L.S., Copeland, E.M., III, et al. (2006). Delayed breast cellulitis: An evolving complication of breast conservation. *International Journal of Radiation Oncology, Biology, Physics*, 66(5), 1339-1346.
- Jeffs, E. (2006). Treating breast cancer-related lymphoedema at the London Haven: Clinical audit results. *European Journal of Oncology Nursing*, 10(1), 71-79.
- Johansson, K., Tibe, K., Weibull, A., & Newton, R.C. (2005). Low intensity resistance exercise for breast cancer patients with arm lymphedema with or without compression sleeve. *Lymphology*, 38(4), 167-180.
- Karadibak, D., Yavuzsen, T., & Saydam, S. (2008). Prospective trial of intensive decongestive physiotherapy for upper extremity lymphedema. *Journal of Surgical Oncology*, 97(7), 572-577.
- Kaviani, A., Fateh, M., Yousefi Nooraie, R., Alinagi-zadeh, M.R., & Ataie-Fashtami, L. (2006). Low-level laser therapy in management of postmastectomy lymphedema. *Lasers in Medical Science*, 21(2), 90-94.
- Koul, R., Dufan, T., Russell, C., Guenther, W., Nugent, Z., Sun, X., et al. (2007). Efficacy of complete decongestive therapy and manual lymphatic drainage on treatment-related lymphedema in breast cancer. *International Journal of Radiation Oncology, Biology, Physics*, 67(3), 841-846.
- Lymphoedema Framework. (2006). *Best practice for the management of lymphoedema*. London: MEP Ltd.
- Mahamaneerat, W.K., Shyu, C.-R., Stewart, B.R., & Armour, J.M. (in press). Post-op swelling and lymphoedema following breast cancer treatment: A baseline-comparison BMI-adjusted approach. *Journal of Lymphoedema*.
- McNeely, M.L., Magee, D.J., Lees, A.W., Bagnall, K.M., Haykowsky, M., & Hanson, J. (2004). The addition of manual lymph drainage to compression therapy for breast cancer related lymphedema: A randomized controlled trial. *Breast Cancer Research and Treatment*, 86(2), 95-106.
- Moffatt, C.J., Franks, P.J., Doherty, D.C., Williams, A.F. (2003). Lymphoedema: An underestimated health problem. *QJM*, 96 (10), 731-738.
- Moseley, A.L., Carati, C.J., & Piller, N.B. (2007). A systematic review of common conservative therapies for arm lymphoedema secondary to breast cancer treatment. *Annals of Oncology*, 18(4), 639-646.

- National Lymphedema Network Medical Advisory Committee. (2004). *Position statement of the National Lymphedema network. Topic: Air travel*. Retrieved January 9, 2008, from <http://www.lymphnet.org/pdfDocs/nlnairtravel.pdf>.
- National Lymphedema Network Medical Advisory Committee (2005). *Position statement of the National Lymphedema Network. Topic: Lymphedema risk reduction practices*. Retrieved January 9, 2008 from <http://www.lymphnet.org/pdfDocs/nlnriskreduction.pdf>.
- National Lymphedema Network. (2008a). *NLN position paper: Air travel*. Retrieved May 15, 2008, from <http://www.lymphnet.org/pdfDocs/nlnairtravel.pdf>
- National Lymphedema Network. (2008b). *NLN position paper: Lymphedema risk reduction practices*. Retrieved May 15, 2008, from <http://www.lymphnet.org/pdfDocs/nlnriskreduction.pdf>
- NHMRC National Breast Cancer Centre, Australia. (1997). Lymphoedema: Prevalence, risk factors and management: A review of research. *The management of lymphoedema*. Retrieved from <http://www.nbcc.org.au/pages/info/resource/nbccpubs/lympho/manage.htm>.
- Ridner, S.H., & Dietrich, M.S. (2008). Self-reported comorbid conditions and medication usage in breast cancer survivors with and without lymphedema. *Oncology Nursing Forum*, 35(1), 57-63.
- Shaw, C., Mortimer, P., & Judd, P.A. (2007). Randomized controlled trial comparing a low-fat diet with a weight-reduction diet in breast cancer-related lymphedema. *Cancer*, 109(10), 1949-1956.
- Soran, A., D'Angelo, G., Begovic, M., Ardic, F., Harlak, A., Samuel Wieand, H., et al. (2006). Breast cancer-related lymphedema: What are the significant predictors and how they affect the severity of lymphedema? *Breast Journal*, 12(6), 536-543.
- Stout Gergich, N.L., Pfalzer, L.A., McGarvey, C., Springer, B., Gerber, L.H., & Soballe, P. (2008). Preoperative assessment enables the early diagnosis and successful treatment of lymphedema. *Cancer*, 112(12), 2809-2819.
- Szuba, A., Achalu, R., & Rockson, S.G. (2002). Decongestive lymphatic therapy for patients with breast carcinoma-associated lymphedema. *Cancer*, 95(11), 2260-2267.
- Vignes, S., Porcher, R., Arrault, M., & Dupuy, A. (2007). Long-term management of breast cancer-related lymphedema after intensive decongestive physiotherapy. *Breast Cancer Research and Treatment*, 101(3), 285-290.
- Williams, A.F., Vadgama, A., Franks, P.J. & Mortimer, P.S. (2002). A randomized controlled crossover study of manual lymphatic drainage therapy in women with breast cancer-related lymphoedema. *European Journal of Cancer Care* 11(3), 254-261.
- Woods, M. (2003). The experience of manual lymph drainage as an aspect of treatment for lymphoedema. *International Journal of Palliative Nursing*, 9(4), 336-342.
- Young-McCaughan, S., & Arzola, S.M. (2007). Exercise intervention research for patients with cancer on treatment. *Seminars in Oncology Nursing*, 23(4), 264-274.