

Nausea and Vomiting

References

- Aapro, M.S., Grunberg, S.M., Manikhas, G.M., Olivares, G., Suarez, T., Tjulandin, S.A., et al. (2006). A phase III, double-blind, randomized trial of palonosetron compared with ondansetron in preventing chemotherapy-induced nausea and vomiting following highly emetogenic chemotherapy. *Annals of Oncology*, 17(9), 1441-1449.
- Abali, H., & Celik, I.S.B.,(2007). Tropisetron, ondansetron, and granisetron for control of chemotherapy-induced emesis in Turkish cancer patients: A comparison of efficacy, side-effect profile, and cost. *Cancer Investigation*, 25(3), 135-139.
- Ahles, T.A., Tope, D.M., Pinkson, B., Walch, S., Hann, D., Whedon, M., et al. (1999). Massage therapy for patients undergoing autologous bone marrow transplantation. *Journal of Pain and Symptom Management*, 18, 157-163.
- Arakawa, S. (1997). Relaxation to reduce nausea, vomiting, and anxiety induced by chemotherapy in Japanese patients. *Cancer Nursing*, 20, 342-349.
- American Society of Health System Pharmacists. (1999). ASHP therapeutic guidelines on the pharmacologic management of nausea and vomiting in adult and pediatric patients receiving chemotherapy or radiation therapy or undergoing surgery. *American Journal of Health System Pharmacy*, 56, 729-764.
- Andersen C. Adamsen L. Moeller T. Midtgaard J. Quist M. Tveteraas A. Rorth M. (2006). The effect of a multidimensional exercise programme on symptoms and side-effects in cancer patients undergoing chemotherapy--the use of semi-structured diaries. *European Journal of Oncology Nursing*, 10(4), 247-262.
- Antiemetic Subcommittee of the Multinational Association of Supportive Care in Cancer. (2006). Prevention of chemotherapy- and radiotherapy-induced emesis: Results of the 2004 Perugia International Antiemetic Consensus Conference. *Annals of Oncology*, 17, 20-28.
- Billhult A, Bergbom I, Stener-Victorin E. (2007). Massage relieves nausea in women with breast cancer who are undergoing chemotherapy. *The Journal of Alternative and Complementary Medicine*, 13(1), 53-57.
- Bleicher, J., Bhaskara, A., Huyck, T., Constantino, S., Bardia, A., Loprinzi, C.L., Silberstein, P.T. (2008). Lorazepam, diphenhydramine, and haloperidol gel for rescue from chemotherapy-induced nausea and vomiting: Results of two pilot trials. *Journal of Supportive Oncology*. 6(1), 27-32.
- Borjesson, S., Hursti, T.J., Tishelman, C., Peterson, C., & Steineck, G. (2002). Treatment of nausea and emesis during cancer chemotherapy: Discrepancies between antiemetic effect and well-being. *Journal of Pain and Symptom Management*, 24, 345-358.
- Booth CM, Clemons M, Dranitsaris G, Joy A, Young S, Callaghan W, Trudeau M, Petrella T. (2007). Chemotherapy-induced nausea and vomiting in breast cancer patients: a prospective observational study. *Journal of Supportive Oncology*. 5(8),374-380.
- Buckle, J. (2007). Complementary therapy. Literature review: Should nursing take aromatherapy more seriously?*British Journal of Nursing* 16 (2), 116-120.
- Cassileth, B.R., & Vickers, A.J. (2004). Massage therapy for symptom control: Outcome study at a major cancer center. *Journal of Pain and Symptom Management*, 28, 244-249.
- Choo SP, Kong KH, Lim WT, Gao F, Chua K, Leong SS. (2006). Electroacupuncture for refractory acute emesis caused by chemotherapy. *The Journal of Alternative and Complementary Medicine*, 12(10), 963-969.
- Collins, K.B., & Thomas, D.J. (2004). Acupuncture and acupressure for the management of chemotherapy-induced nausea and vomiting. *Journal of the American Academy of Nurse Practitioners*, 16 (2), 76-80.

Nausea and Vomiting

References

- Davis, M.P. (2008). Oral nabilone capsules in the treatment of chemotherapy-induced nausea and vomiting and pain. *Expert Opinion on Investigational Drugs*, 17(1), 85-95.
- de Carvalho EC, Martins FTM, dos Santos CB. (2007). A pilot study of a relaxation technique for management of nausea and vomiting in patients receiving cancer chemotherapy. *Cancer Nursing*, 30(2), 163-167.
- Dibble, S.L., Chapman, J., Mack, K.A., & Shih, A. (2000). Acupressure for nausea: Results of a pilot study. *Oncology Nursing Forum*, 27, 41-47.
- Dibble SL, Luce J, Cooper BA, Israel J, Cohen M, Nussey B, Rugo H. (2007). Acupressure for chemotherapy-induced nausea and vomiting: a randomized clinical trial. *Oncology Nursing Forum*. 34(4):813-820.
- Einhorn LH, Brames MJ, Dreicer R, Nichols CR, Cullen MT Jr, Bubalo J. (2007). Palonosetron plus dexamethasone for prevention of chemotherapy-induced nausea and vomiting in patients receiving multiple-day cisplatin chemotherapy for germ cell cancer. *Supportive Care in Cancer*, 15(11),1293-1300.
- Ernst, E., & Pittler, M.H. (2000). Efficacy of ginger for nausea and vomiting: A systematic review of randomized clinical trials. *British Journal of Anaesthesia*, 84, 367-71.
- Ezzo, J., Vickers, A., Richardson, M.A., Allen, C., Dibble, S.L., Issell, B. et al. (2005). Acupuncture-point stimulation for chemotherapy-induced nausea and vomiting. *Journal of Clinical Oncology*, 23, 7188-7198.
- Ezzone, S., Baker, C., Rosselet, R., & Terepka, E. (1998). Music as an adjunct to antiemetic therapy. *Oncology Nursing Forum*, 25, 1551-1556.
- Fellowes, D., Barnes, K., & Wilkinson, S. (2004). Aromatherapy and massage for symptom relief in patients with cancer. *Cochrane Database of Systematic Reviews*, 2, CD002287.
- Gamaluddin, S., Kirchner, J.T. (2006). Acupuncture: What the evidence shows now. *Family Practice Recertification* 28 (9), 26-28,33-37.
- Gardani, G., Cerrone, R., Biella, C., Galbiati, G., Proserpio, E., Casiraghi, M., et al. (2007). A progress study of 100 cancer patients treated by acupressure for chemotherapy-induced vomiting after failure of pharmacological approach. *Minerva Medica*, 98(6), 665-668.
- Gralla, R.J., Osoba, D., Kris, M.G., Kirkbride, P., Hesketh, P.J., Chinnery, L.W., et al. (1999). Recommendations for the use of antiemetics: Evidence-based, clinical practice guidelines. *Journal of Clinical Oncology*, 17, 2971-2994.
- Grealish, L., Lomasney, A., & Whiteman, B. (2000). Foot massage: A nursing intervention to modify the distressing symptoms of pain and nausea in patients hospitalized with cancer. *Cancer Nursing*, 23, 237-243.
- Gurpide A, Sadaba B, Martin-Algarra S, Azanza JR, Lopez-Picazo JM, Campanero MA, Cabello JP, Gil-Aldea I, de la Cruz S, Fernandez Gallego V, Reyna C, Olier Garate C, Blanco-Prieto MJ, Ceballos J, Garcia-Foncillas J, Perez-Gracia JL. (2007). Randomized crossover pharmacokinetic evaluation of subcutaneous versus intravenous granisetron in cancer patients treated with platinum-based chemotherapy. *Oncologist*, 12(9),1151-1155.
- Hajdenberg J, Grote T, Yee L, Arevalo-Araujo R, Latimer LA. (2006). Infusion of palonosetron plus dexamethasone for the prevention of chemotherapy-induced nausea and vomiting. *Journal of Supportive Oncology*, 4(9),467-471.
- Herrington, J.D., Jaskiewicz, A.D., & Song, J. (2008). Randomized, placebo-controlled, pilot

Nausea and Vomiting

References

- study evaluating aprepitant single dose plus palonosetron and dexamethasone for the prevention of acute and delayed chemotherapy-induced nausea and vomiting. *Cancer*, 112(9), 2080-2087.
- Herrstedt, J., Roila, F., & ESMO Guidelines Working Group. (2008). Chemotherapy-induced nausea and vomiting: ESMO clinical recommendations for prophylaxis. *Annals of Oncology*, 19 (Suppl. 2), ii110-ii112.
- Herrstedt, J., Sigsgaard, T.C., Nielsen, H.A., Handberg, J., Langer, S.W., Ottesen, S. & Dombernowsky, P. (2007). Randomized, double-blind trial comparing the antiemetic effect of tropisetron plus metopimazine with tropisetron plus placebo in patients receiving multiple cycles of multiple-day cisplatin-based chemotherapy. *Supportive Care in Cancer*, 15 (4), 417-426.
- Hesketh, P.J., Grunberg, S.M., Herrstedt, J., de Wit, R., Gralla, R.J., Carides, A.D., et al. (2006). Combined data from two phase III trials of the NK1 antagonist aprepitant plus a 5HT₃ antagonist and a corticosteroid for prevention of chemotherapy-induced nausea and vomiting: Effect of gender on treatment response. *Supportive Care in Cancer*, 14(4), 354-360.
- Keegan, G.T. & Keegan, L. (2006). The use of ginger to alleviate nausea. *Alternative Therapies in Women's Health*, 8 (8), 57-61.
- Khamales, S., Bethune-Volters, A., Chidiac, J., Bensaoula, O., Delgado, A., & Di Palma, M.. (2006). A randomized, double-blind trial assessing the efficacy and safety of sublingual metopimazine and ondansetron in the prophylaxis of chemotherapy-induced delayed emesis. *Anticancer Drugs*, 17(2), 217-224.
- Kim, S., Shin, I., Kim, J., Kim, Y., Kim, K, Kim, Ki, Yang, S., Yoon, J. (2008). Effectiveness of mirtazapine for nausea and insomnia in cancer patients with depression. *Psychiatry and Clinical Neurosciences*. 62, 75-83.
- King, C.R. (1997). Nonpharmacologic management of chemotherapy-induced nausea and vomiting. *Oncology Nursing Forum*, 24 (7, Suppl.), 41-48.
- Klein, J., & Griffiths, P. (2004). Acupressure for nausea and vomiting in cancer patients receiving chemotherapy. *British Journal of Community Nursing*, 9, 383-387.
- Kris, M.G., Hesketh, P.J., Somerfield, M.R., Feyer, P., Clark-Snow, R., et al. (2006). American Society of Clinical Oncology guideline for antiemetics in oncology: Update 2006. *Journal of Clinical Oncology*, 24(18), 2932-2947.
- Lajolo PP, del Giglio A. (2007). Skipping day 2 antiemetic medications may improve chemotherapy induced delayed nausea and vomiting control: results of two pilot phase II trials. *Supportive Care in Cancer*, 15(3),343-346.
- Luebbert, K., Dahme, B., & Hasenbring, M. (2001). The effectiveness of relaxation training in reducing treatment-related symptoms and improving emotional adjustment in acute non-surgical cancer treatment: A meta-analytical review. *Psycho-Oncology*, 10, 490-502.
- Manusirivithaya, S., Sripramote, M., Tangjitgamol, S., Sheanakul, C., Leelahakorn, S., Thavaramara, T., et al. (2004). Antiemetic effect of ginger in gynecologic oncology patients receiving cisplatin. *International Journal of Gynecologic Cancer*, 14, 1063-1069.
- Marchioro, G., Azzarello, G., Viviani, F., Barbato, F., Pavanetto, M., Rosetti, F., et al. (2000). Hypnosis in the treatment of anticipatory nausea and vomiting in patients receiving cancer chemotherapy. *Oncology*, 59, 100-104.
- Mayer, D.J. (2000). Acupuncture: An evidence-based review of the clinical literature. *Annual Review of Medicine*, 51, 49-63.

Nausea and Vomiting

References

- Meiri, E., Jhangiani, H., Vredenburgh, J.J., Barbato, L.M., Carter, F.J., Yang, H.M., et al. (2007). Efficacy of dronabinol alone and in combination with ondansetron versus ondansetron alone for delayed chemotherapy-induced nausea and vomiting. *Current Medical Research and Opinion*, 23(3), 533-543.
- Melchart D, Ihbe-Heffinger A, Leps B, von Schilling C, Linde K. (2006). Acupuncture and acupressure for the prevention of chemotherapy-induced nausea--a randomised cross-over pilot study. *Supportive Care in Cancer*. 14(8),878-882.
- Miller, M., & Kearney, N. (2004). Chemotherapy-related nausea and vomiting-Past reflections, present practice and future management. *European Journal of Cancer Care*, 13 (1), 71-81.
- Mok TS, Yeo W, Johnson PJ, Hui P, Ho WM, Lam KC, Xu M, Chak K, Chan A, Wong H, Mo F, Zee B. (2007). A double-blind placebo-controlled randomized study of Chinese herbal medicine as complementary therapy for reduction of chemotherapy-induced toxicity. *Annals of Oncology*, 18(4), 768-774.
- Molassiotis A, Helin AM, Dabbour R, Hummerston S. (2007). The effects of P6 acupressure in the prophylaxis of chemotherapy-related nausea and vomiting in breast cancer patients. *Complementary Therapies in Medicine*, 15(1),3-12.
- Molassiotis, A., Yung, H.P, Yam, B.M.C., Chan, F.Y.S., & Mok, T.S.K. (2002). The effectiveness of progressive muscle relaxation training in managing chemotherapy-induced nausea and vomiting in Chinese breast cancer patients: A randomised controlled trial. *Supportive Care in Cancer*, 10, 237-246.
- Multinational Association of Supportive Care in Cancer. (2008). Antiemetic guidelines. Retrieved June 23, 2008 from <http://www.mascc.org>
- National Comprehensive Cancer Network. (2006). Antiemesis practice guidelines in oncology, version 1.2006. Retrieved March 21, 2006 from <http://www.nccn.org>.
- National Comprehensive Cancer Network. (2008). Antiemesis practice guidelines in oncology, version 2.2008. Retrieved March 14, 2008 from <http://www.nccn.org>.
- Navari RM, Einhorn LH, Loehrer PJ Sr, Passik SD, Vinson J, McClean J, Chowhan N, Hanna NH, Johnson CS. (2007). A phase II trial of olanzapine, dexamethasone, and palonosetron for the prevention of chemotherapy-induced nausea and vomiting: a Hoosier oncology group study. *Supportive Care in Cancer*, 15(11),1285-1291.
- Oyama, H., Kaneda, M., Katsumata, N., Akechi, T., & Ohsuga, M. (2000). Using the bedside wellness system during chemotherapy decreases fatigue and emesis in cancer patients. *Journal of Medical Systems*, 24 (3), 173-182.
- Pectasides, D., Dafni, U., Aravantinos, G., Timotheadou, E., Skarlos, D.V., Pavlidis, N., et al. (2007). A randomized trial to compare the efficacy and safety of antiemetic treatment with ondansetron and ondansetron zydis in patients with breast cancer treated with high-dose epirubicin. *Anticancer Research*, 27(6C), 4411–4417.
- Polovich, M., White, J.M., & Kelleher, L.O. (Eds.). (2005). *Chemotherapy and biotherapy guidelines and recommendations for practice* (2nd ed.). Pittsburgh, PA: Oncology Nursing Society.
- Raghavendra, R.M., Nagarathna, R., Nagendra, H.R., Gopinath, KS, Srinath, B.S., Ravi, B.D., Patil, S., Ramesh, B.S., Nalini, R. (2007). Effects of an integrated yoga programme on chemotherapy-induced nausea and emesis in breast cancer patients. *European Journal of Cancer Care*, 16, 462-474.

Nausea and Vomiting

References

- Redd, W.H., Montgomer, G.H. & DuHamel, K.N. (2001). Behavioral intervention for cancer treatment. *Journal of the National Cancer Institute*, 93, 810-823.
- Richardson J, Smith JE, McCall G, Richardson A, Pilkington K, Kirsch I. (2007). Hypnosis for nausea and vomiting in cancer chemotherapy: a systematic review of the research evidence. *European Journal of Cancer Care*, 16(5),402-412.
- Roscoe, J.A., Matteson, S.E., Morrow, G.R., Hickok, J.T., Bushunow, P., Griggs, J., et al.(2005). Acustimulation wrist bands are not effective for the control of chemotherapy-induced nausea in women with breast cancer. *Journal of Pain and Symptom Management*, 29, 376-384.
- Roscoe, J.A., Morrow, G.R., Hickok, J.T., Bushunow, P., Pierce, I., Flynn, P.J., et al. (2003). The efficacy of acupressure and acustimulation wrist bands for the relief of chemotherapy-induced nausea and vomiting: A University of Rochester Cancer Center Community Clinical Oncology Program multicenter study. *Journal of Pain and Symptom Management*, 26, 731-740.
- Roscoe, J.A., Morrow, G.R., Matteson, S., Bushunow, P., & Tian, L. (2002). Acustimulation wristbands for the relief of chemotherapy-induced nausea. *Alternative Therapies*, 8 (4), 56-63.
- Sahler, O.J.Z, Hunter, B.C, & Liesveld, J.L. (2003). The effect of using music therapy with relaxation imagery in the management of patients undergoing bone marrow transplantation: A pilot feasibility study. *Alternative Therapies*, 9 (6), 70-74.
- Schmoll, H.J., Apro, M.S., Poli-Bigelli, S., Kim, H.K., Park, K., Jordan, K., et al. (2006). Comparison of an aprepitant regimen with a multiple-day ondansetron regimen, both with dexamethasone, for antiemetic efficacy in high-dose cisplatin treatment. *Annals of Oncology*, 17(6), 1000-1006.
- Shen, J., Wenger, N., Glaspy, J., Hays, R.D., Albert, P.S., Choi, C., et al. (2000). Electroacupuncture for control of myeloablative chemotherapy-induced emesis: A randomized controlled trial. *JAMA*, 284, 2755-2761.
- Shi, Y., He, X., Yang, S., Ai, B., Zhang, C., Huang, D., et al. (2007). Ramosetron versus ondansetron in the prevention of chemotherapy-induced gastrointestinal side effects: A prospective randomized controlled study. *Chemotherapy*, 53(1), 44-50.
- Shin, Y.H., Kim, T.I., Shin, M.S., & Juon, H. (2004). Effect of acupressure on nausea and vomiting during chemotherapy cycle for Korean postoperative stomach cancer patients. *Cancer Nursing*, 27, 267-274.
- Taixiang W. Munro AJ. Guanjian L. (2005). Chinese medical herbs for chemotherapy side effects in colorectal cancer patients. *Cochrane Database of Systematic Reviews*, 1 CD004540.
- Treish, I., Shord, S., Valgus, J., Harvey, D., Nagy, J., Stegal, J., et al. (2003). Randomized double-blind study of the Reliefband as an adjunct to standard antiemetics in patients receiving moderately-high to highly emetogenic chemotherapy. *Supportive Care in Cancer*, 11, 516-521.
- Troesch, L.M., Rodehaver, C.B., Delaney, E.A., & Yanes, B. (1993). The influence of guided imagery on chemotherapy-related nausea and vomiting. *Oncology Nursing Forum*, 20, 1179-1185.
- Vrabel, M. (2007). Is ondansetron more effective than granisetron for chemotherapy-induced nausea and vomiting? A review of comparative trials. *Clinical Journal of Oncology Nursing*, 11(6), 809-813.

Nausea and Vomiting

References

- Williams, S.A., & Schreier, A.M. (2004). The effect of education in managing side effects in women receiving chemotherapy for treatment of breast cancer [Online exclusive]. *Oncology Nursing Forum*, 31, E16-E23.
- Winningham, M.L., & MacVicar, M.G. (1988). The effect of aerobic exercise on patient reports of nausea. *Oncology Nursing Forum*, 15, 447-450.
- Zhang M. Liu X. Li J. He L. Tripathy D. (2007). Chinese medicinal herbs to treat the side-effects of chemotherapy in breast cancer patients. *Cochrane Database of Systematic Reviews*, 3, CD004921.