



# Evaluation of a Preoperative Clinic for Women With Gynecologic Cancer

Fang Huang, DNP, RN, Yen Y. Chia, PhD, RN, Chui L. Eng, MSN, RN, Yong K. Lim, MBBS, MRCOG, Kwai L. Yam, MBBS, FRCOG, Soh C. Tan, MSN, RN, and Marilyn Hockenberry, PhD, RN, PPCNP-BC, FAAN

Although many patients with gynecologic cancer undergo surgery, time constraints during the preoperative consultation may affect the accuracy of the information exchange, as well as compromise the quality of the patient assessment and care plan. Both put patients at a higher risk for complications during surgery and the postoperative period. This article describes an advanced practice RN-led preoperative assessment and education clinic designed to improve the quality of preoperative preparation and postoperative outcomes of patients with gynecologic cancer.

## At a Glance

- Among patients with gynecologic cancer, surgery is the major treatment modality.
- The success of surgery depends on many factors, and a failure to identify those risks may negatively affect surgical outcomes and hospital length of stay.
- A preoperative assessment and education clinic led by advanced practice RNs has been shown to improve preoperative preparation and postoperative outcomes in patients with gynecologic cancer.

Fang Huang, DNP, RN, is an advanced practice nurse, Yen Y. Chia, PhD, RN, is a deputy director of nursing, and Chui L. Eng, MSN, RN, is an assistant director of nursing and an advanced practice nurse, all in the Division of Nursing at the KK Women's and Children's Hospital in Singapore; Yong K. Lim, MBBS, MRCOG, is a senior consultant and the department head and Kwai L. Yam, MBBS, FRCOG, is a senior consultant, both in the Gynaecological Oncology Department at the KK Women's and Children's Hospital; Soh C. Tan, MSN, RN, is the chief nursing officer at the Ministry of Health in Singapore; and Marilyn Hockenberry, PhD, RN, PPCNP-BC, FAAN, is a professor in the School of Nursing at Duke University in Durham, NC. The authors take full responsibility for the content of the article. This project was supported, in part, by a grant from the KKH Health Endowment Fund. No financial disclosures relevant to the content of this article have been disclosed by the editorial staff. Huang can be reached at [huangfangjm@hotmail.com](mailto:huangfangjm@hotmail.com), with copy to editor at [CJONEditor@ons.org](mailto:CJONEditor@ons.org).

Key words: gynecologic cancer; preoperative assessment; preoperative preparation; nurse-led clinic  
Digital Object Identifier: 10.1188/15.CJON.769-772

The burden of cancer diagnosis and treatment for women with gynecologic cancer is profound. Surgical intervention is the major treatment modality for this patient population, but the success of surgical interventions depends on a complex interplay of factors, including the patient's physical condition, psychological distress, financial stability, and social support. Failure to identify those risk factors in women with gynecologic cancer prior to surgery has been shown to adversely affect surgical outcomes

and hospital length of stay (LOS) (Dean, Finan, & Kline, 2001; Ellis, Spiers, Coutts, Fairburn, & McCracken, 2012; Fleisher, 2009; Vilar-Compte et al., 2008).

Increases in surgical complications and psychological distress in preoperative patients with gynecologic cancer have been attributed to inadequate preoperative assessment, preparation, and planning, as well as limited opportunities for clinic staff to increase patients' understanding of the diagnosis and recommended treatment (Cimprich, 1999; Love, 2004; Mc-

Corkle et al., 2009). Ideally, an individual patient risk assessment and tailored preoperative preparation plan should be created in the outpatient setting at the initial cancer diagnosis (Schofield et al., 2003). Preoperative patient assessments are known to lower patient anxiety while increasing identification of modifiable risk factors for surgical complications. However, a drastic rise in patient volume and an increase in patient and family expectations for additional information related to the diagnosis and care plan have placed time constraints on the healthcare team, resulting in unmet expectations and compromised information exchange. This, in turn, puts patients at a higher risk of complications during the perioperative period (Jie et al., 2012; Seibaek, Blaakaer, Petersen, & Hounsgaard, 2013). Although the importance of preoperative information is well recognized, studies continuously report that patients lack adequate information, care coordination, and therapeutic relationships with providers (Guest, Manderville, & Thomson, 2012; Jolley, 2007).

## Background

Clinics led by advanced practice RNs (APRNs) have been developed to address time constraints and fulfill patients' perceived unmet needs, identify risks, improve the quality of service, and address actual or potential problems through teamwork (Loftus & Weston, 2001). For example, evaluation of a nurse-coordinated multidisciplinary preoperative assessment program for older adults demonstrated a reduction in LOS from 8.9 to 4.9 days and a decrease in serious postoperative complications from about 9% to 2% (Ellis